

Magnetosphere response to magnetic clouds: multi-satellite observations during 1995-1998

Yu. I. Yermolaev¹, G. N. Zastenker¹, N. S. Nikolaeva¹, J.-A. Sauvaud², K.W.Ogilvie³

¹ Space Research Institute, Russian Academy of Sciences, Profsoyuznaya 84/32, 117810 Moscow, Russia

² CESR/CNES, Toulouse, France

³ NASA/GSFC, Greenbelt, USA

Received: 31 December 2000 / Revised: 12 May 1997 / Accepted: 6 June 1997

Abstract. On the basis of ISTP spacecraft and ground observations during first 40 months of INTERBALL operation in 1995-1998 we study magnetosphere response to magnetic cloud passages including geomagnetic storms and polar activations. During this time 35 magnetic clouds were measured in the solar wind which resulted in 15 from 19 strong (peak $D_{st} < -100$ nT) magnetic storms observed on the ground stations. The low and moderately high changes in magnetic cloud IMF and solar wind parameter variations result in usual magnetosphere response to the similar changes without magnetic cloud passages. Extremely high jumps of parameters in the magnetic clouds generate unusual response: (1) strong and complicated magnetosphere compression and deformation relative to average locations; (2) large-amplitude oscillations of geomagnetic tail structures past satellites, and (3) acceleration of ions and electrons in the plasma sheet and their injections in the polar regions. During magnetic clouds the value of peak $|D_{st}|$ correlates with numbers of polar activations, and the same dependence is observed for strong magnetic storms.

Key words. Magnetic cloud, magnetosphere, magnetic storms and substorms

1 Introduction

One of the main problems of the solar-terrestrial physics is a question which magnetosphere responses are caused by different variations in solar/interplanetary medium. There are two aspects. On the one hand, the problem plays key role in our understanding of geophysics. On another hand, this knowledge presents a practical interest in many areas of mankind activity.

The large group of papers describes the processes of solar wind energy input into magnetosphere and development

of magnetospheric disturbances (see, e.g. (Gonzales *et al.* (1994, 1999); Kamide *et al.* (1998); Petrukovich and Klimov (2000); Wilson (2000) and references therein). It was shown that the existence of southward component of the interplanetary magnetic field (IMF) results in input of solar wind energy into magnetosphere and its accumulation in the magnetic tail. Then when this energy reaches sufficient level it can begin to release as reconfiguration of current systems and as plasma acceleration or heating, which results in the magnetosphere disturbances, such as magnetic storms and substorms.

Another group of investigators studied selected events in the solar atmosphere, in the interplanetary space and in the magnetosphere and correlations between them (see, e.g. Gosling *et al.* (1991); Webb (1995); Tsurutani *et al.* (1995); Crooker (2000) and references therein). They found that geoeffective events (in the sense that they can cause geomagnetic storms) are mainly the magnetic clouds (MC), which are interplanetary manifestation of the coronal mass ejections, and corotating interaction regions (CIR) derived from interaction of fast and slow streams of solar wind. MCs and CIRs are geoeffective because they are faster than ambient plasma and compress any southward IMF in the vicinity of their edges or inside of event body. Correlation with polar magnetospheric disturbances is not sufficiently studied yet because these disturbances have characteristic time of several tens minutes and they should be compared not with the magnetic cloud as a whole, but with its separate structures and disturbances, such as the interplanetary shock (IS) before MC, the leading and trailing edges (LE and TE) of MC, the IS before TE, the jump of plasma pressure, the changes in IMF magnitude and orientation. It is also important to study displacement of magnetosphere boundaries (including the bow shock and magnetopause) under these unusual interplanetary conditions. Such an analysis for several strong magnetic clouds has already been done on the basis of multi-satellite INTERBALL project (Yermolaev *et al.*, 1997a, 1998, 2000a,b), and here we summarize the results of our analysis on the basis of full statistics of magnetic clouds during first 40 months of INTERBALL observations (August, 1995 - December, 1998). We limited our study by time interval of relatively low solar activity before maximum of solar cycle.

2 General view of interval

It has been known that substantial variations occur over the 11-year solar cycle in disturbances of solar wind and Earth's magnetosphere. We study time interval in vicinity of minimum (1996) and growth phase of cycle. To evaluate magnetosphere disturbances the geomagnetic indices measured at the ground stations are usually used. The D_{st} index, which connects with the geomagnetic field near the equator and disturbance of the ring current, describes well the development of the global large-scale disturbances - magnetic storms.

Figures 1 - 3 show hour-averaged values of D_{st} index (<http://spidr.ngdc.noaa.gov>) during 47 Sun's rotations (40 months from August, 1995 to December, 1998). Large magnetic storms (peak $D_{st} \leq -100$ nT) are indicated by red triangles. There were 19 large storms, and their number slightly increased in the end of interval, the closer to maximum of solar cycle. This tendency is confirmed by D_{st} index data for 1999-2000 period when there were 17 strong magnetic storms during 24 months.

To analyze interplanetary conditions for magnetic storms we use the key parameters of plasma and magnetic field measured by WIND Ogilvie *et al.* (1995); Lepping *et al.* (1995) and, in some cases, by the other spacecraft (SOHO and IMP-8) (<http://cdaweb.gsfc.nasa.gov>). Green and blue horizontal lines in top of pannels present time intervals of MC and CIR observations, respectively, and red vertical lines IS preceding them. Characteristic behaviour of plasma and magnetic field in MC and CIR has been early discussed in the literature and may be found in papers by Gosling and Pizzo (1999) and Crooker (2000). Figures 1 - 3 show that the 19 large storms were connected with 15 magnetic clouds and 4 corotating interaction regions. At the same time the analysis of data indicated at least 35 MCs during August, 1995 - December, 1998. A part of them was studied earlier (Yermolaev *et al.*, 1998, 2000a,b), another part was added from the list of coronal mass ejections (Gopalswamy *et al.*, 2000), and a part was selected recently. Thus we have possibility to compare MCs generated different magnetic storms in wide range of D_{st} .

The list of considered events is presneted in Table 1, which includes the date and duration of MC observations (the interval between MC IS and LE is indicated additionally in brackets). Also indicated here are the regions of space where was crossed at this time by INTERBALL/Tail Probe (INTERBALL-1 hereafter) satellite: SW is the solar wind, MSH - the magnetosheath, MS - magnetosphere (the tail lobes, plasma and neutral sheets, mantle, LLBL and PSBL). As seen from Table 1, INTERBALL-1 in the majority cases was in different regions of the magnetosphere and measured the parameters of plasma, magnetic field and energetic particles there. The INTERBALL/Auroral Probe (INTERBALL-2) satellite with a low-apogee 6-hour orbit measured various parameters in the polar magnetosphere. Owing to a variety of satellite locations at the time MC passages, we have a possibility of investigating the different magnetosphere region under different solar wind conditions.




Fig. 3. One-hour averaged D_{st} variations for June - December, 1998

3 Geoeffectiveness of magnetic clouds

As indicated in Figures 1 - 3, the magnetic storm durations are close to magnetic cloud ones. For instance for January 10-11, 1997 magnetic cloud (Burlaga *et al.*, 1998), duration of magnetic storm and magnetic cloud were ~ 18 h and 23 h, respectively. Therefore, we can compare the instant of magnetic storm beginning with MC structure. On the whole, the large drop of D_{st} index is observed after southward IMF turning with 0-2 h delay (We used 1-hour averaged D_{st} index data). Usually these IMF turnings occurred in compressed region between IS and MC LE or inside of MC body due to slow IMF rotation here.

To describe the polar disturbances we used either the magnetic field data of several polar ground stations, near which the events took place, or the integral polar indices.




Fig. 1. One-hour averaged D_{st} variations for August, 1995 - December, 1996




Fig. 2. One-hour averaged D_{st} variations for January, 1997 - May, 1998

Table 1. Magnetic clouds observed on INTERBALL-1 as well as on WIND (*) and SOHO+IMP-8 (**).

| N | Date | Durations, h | | Space regions by | Conditions in SW |
|---------|---------------|--------------|-----------|------------------|--------------------------------------|
| | | MC | (+ Shock) | INTERBALL-1 | |
| 1995 | | | | | |
| 1 | Aug.22-23 * | 19 | (+7) | MS/MSH/SW | B_z and P jumps |
| 2 | Oct.18-19 * | 28 | (+8) | MS/MSH/MS | B_z variations |
| 1996 | | | | | |
| 3 | Dec.24-25 * | 33 | (+10) | MS/MSH/MS | P jumps |
| 1997 | | | | | |
| 4 | Jan.10-11 * | 23 | (+4) | MS/MSH/MS | SW with $N \sim 150 \text{ cm}^{-3}$ |
| 5 | Feb. 9-11 * | 41 | (+14) | SW/MSH/MS/MSH/SW | $B_z < 0$ |
| 6 | Apr.10-11 * | 22 | (+9) | SW | Sharp B_z changes |
| 7 | Apr. 21-23 * | 43 | (+1) | SW | P jumps |
| 8 | May 15-16 * | 46 | (+4) | SW | B_z changes |
| 9 | June 8-9 * | 24 | (+3) | MS/MSH/SW | Multiple P jumps |
| 10 | June 19 * | 10 | (+6) | SW | Quiet SW |
| 11 | July 15-16 * | 45 | (+6) | SW | $B_z < 0$ |
| 12 | Aug. 3-4 * | 13 | (+4) | SW/MSH/SW | Quiet SW |
| 13 | Sept. 3 * | 12 | (+10) | SW/MSH/MS | B_z jumps |
| 14 | Sept.18-20 * | 56 | (+4) | MS/MSH/MS | B_z jumps |
| 15 | Sept. 21 * | 5 | (+5) | MSH | P jumps |
| 16 | Sept.21-22 * | 19 | (+3) | MSH/SW | P jumps |
| 17 | Oct. 1-2 ** | 42 | (+4) | MS/MSH/MS | - |
| 18 | Oct. 10-12 * | 45 | (+5) | MSH | P jumps |
| 19 | Nov. 7-8 * | 24 | (+7) | MS | P and B_z variations |
| 20 | Nov. 22-23 * | 18 | (+10) | MS | $B_z < 0$ |
| 21 | Dec. 10-11 * | 15 | (+16) | MS | B_z jumps |
| 22 | Dec. 30-31 * | 25 | (+7) | MS | P and B_z jumps |
| 1998 | | | | | |
| 23 | Jan. 7-8 * | 29 | (+14) | MS | B_z jumps |
| 24 | Feb. 4-5 * | 41 | (+17) | MSH/MS | P jumps |
| 25 | Feb. 17-18 * | 14 | (+16) | MS/MSH | P and B_z jumps |
| 26 | Mar 4-5 * | 30 | (+4) | MS/MSH/SW | B_z jumps |
| 27 | May 2-3 *,** | ? | (+14) | SW | $B_z < 0$ |
| 28 | May 4-5 * | 15 | (+9) | SW/MSH | B_z jumps |
| 29 | June 2 *,** | 8 | ? | SW | Quiet SW |
| 30 | June 24-25 * | 35 | (+4) | SW | P jumps |
| 31 | Sept. 25-26 * | 29 | (+7) | MSH/MS | P jumps |
| 32 | Oct. 18-20 * | 22 | (+9) | MC/MSH | B_z jumps |
| 33 | Nov. 7-8 *,** | ? | (+4) | MS | P and B_z variations |
| 34 | Nov. 8-10 * | 34 | (+13) | MS | P jumps |
| 35 | Nov. 13-14 * | 32 | (+4) | MS | B_z and P jumps |
| Average | | 27 | (+8) | | |

Particularly we analyzed *Contracted Oval*, *Standard Oval*, and *Expanded Oval* calculated for 3 systems of stations located on 3 concentric circles near the northern magnetic pole (For more details see the Auroral Oval Indices on the Cluster/Ground-Based Data Center web site <http://www.wdc.rl.ac.uk/ovals/plots>). The analysis of additional data indicates that these indices are sensitive to substorms and allow us to select them. However, in a small number of cases they demonstrate polar activations which are not substorms. In our analysis we call all these phenomena "activations" and have in mind that about 2/3 of the cases relate

to substorms. Figures 4 - 5 show these indices for magnetic cloud of January 10-12,1997. The comparison of fig.3 in paper by Burlaga *et al.* (1998) and Figs. 4,5 shows that the drop in indices are observed soon after the passage of IS and MC LE. However, the changes in the IMF orientation and jumps in the field magnitude and SW pressure can be found not for all activations. This comparison of polar indices variations with MC structures was made for all MCs shown in Figs. 1 - 3 and Table 1.

Table 2 presents the minimum of hour-averaged values of D_{st} index and number of activations of polar indices re-




Fig. 4. Time dependences of the auroral indices *Contracted Oval*, *Standard Oval*, and *Expanded Oval* on January 10, 1997.

lated to the set of MC structures. For event of January 10-11, 1997, both IS and MC LE correspond to activations (this is designated as 1 activation per 1 structure, i.e. 1/1), while no activation corresponds to MC TE on January 11 (this is designated as 0/1). The decreases of the D_{st} index were observed usually for all MCs. However, in some cases (for example, on September 21, 1997 and on June 24-25, 1998) the index pointed to very weak magnetic storms or even their full absence (on June 2, 1998). In last cases the IMF B_z component was less than -5 nT only during short time intervals (less 1.5 h).

The comparison of activations with the MC structure has shown that only 185 of 237 activations (78% of their total number) can be associated with IS before MC LE (IS1), LE, TE, IS before TE (IS2), the sign of IMF B_z ($B_z < 0$), and the jumps of field (ΔB_z) and dynamics pressure of SW plasma

(ΔP). In this case, the highest relative frequency of activations (the ratio of the number of activations to the number of events of selected type) is observed after IS1 and LE. However, some strong jumps of P and IMF (as, for example, a very high pressure at MC TE on January 11, 1997) have not resulted in activations. It should be noted that the large jump was observed after 8 h of positive IMF B_z .

The data about D_{st} and number of activations presented in Table 2 were shown by black circles in figure 6 and approximated by solid line. The dependence of number of activations NA on D_{st} index for large magnetic storms ($D_{st} < -90$ nT) presented by open circles and dashed line are very closed to dependence for all magnetic clouds. These data allow us to suggest that there is a relation between number of activations NA and D_{st} index. Despite a high scatter of the data at a rather scarce statistics, it can be noted that the passage




Fig. 5. Time dependences of the auroral indices *Contracted Oval*, *Standard Oval*, and *Expanded Oval* on January 11, 1997.

of MC causing strong decrease of D_{st} index is accompanied by a higher number of polar activations (the linear approximation gives the dependence for number of activations $NA = -0.06D_{st} + 1.5$). The problem of relation between slowly varying global geomagnetic indices and rapidly varying polar indices has been already discussed in the literature (see, e.g. Kamide *et al.* (1998) and Wilson (2000)). However, the mechanisms of such a relation during MC passage periods can have some peculiarities and call further investigations.

4 Magnetosphere boundaries

Since the location of the magnetopause (MP) is determined by the balance of plasma and magnetic field pressures in the

solar wind, decelerated and heated at the bow shock (BS), and inside the magnetosphere, any change of conditions in the interplanetary medium results in a displacement of the MP and hence in the displacement of the BS, for which MP is an obstacle when the solar wind flows around it. The INTERBALL-1 satellite locations at BS and MP crossings allow the BS and MP locations to be compared with the solar wind conditions determined by other spacecraft and with model predictions.

We considered 44 MP crossings by the INTERBALL-1 satellite at MC passage time. Figure 7 present the locations of these crossings in the meridional plane (XR_{YZ} , where $R_{YZ} = \sqrt{Y^2 + Z^2}$) and at the cross-section of the tail (YZ), as well as the average locations of MP and BS. It is seen from the figure the deviation of a real MP location from average one (observed at SW pressure of $P \sim 2$ nPa) varies from 1-2

Table 2. Geoeffectiveness of magnetic clouds' structures.

| Date | Dst, nT | Number of substorms and activations | | | | | | | |
|-------------|---------|-------------------------------------|-------|-------|---------------------|-------------------|-------|-----|------|
| | | Total | IS1 | LE | B _z jump | B _z <0 | Pjump | IS2 | TE |
| 1995 | | | | | | | | | |
| Aug. 22-23 | -61 | 3 | 0/1 | 0/1 | 3/5 | 0 | 0/2 | 0/0 | 0/1 |
| Oct. 18-19 | -127 | 9 | 1/1 | 1/1 | 1/1 | 2 | 1/3 | 1/1 | 0/1 |
| 1996 | | | | | | | | | |
| Dec. 23-25 | -33 | 2 | 0/1 | 0/1 | 0/5 | 1 | 1/1 | 0/0 | 0/1 |
| 1997 | | | | | | | | | |
| Jan. 10-11 | -78 | 9 | 0/1 | 1/1 | 2/3 | 2 | 1/3 | 0/1 | 0/1 |
| Feb. 8-11 | -68 | 15 | 1/1 | 1/1 | 2/4 | 3 | 3/5 | 0/0 | 1/1 |
| Apr. 10-11 | -82 | 4 | 0/1 | 1/1 | 1/2 | 0 | 0/0 | 0/0 | 1/1 |
| Apr. 21-23 | -107 | 2 | 1/1 | 0/1 | 0/5 | 0 | 1/1 | 0/0 | 0/0 |
| May 15-16 | -115 | 6 | 0/1 | 1/1 | 2/6 | 3 | 0/1 | 0/0 | 1/1 |
| June 8-9 | -84 | 5 | 1/1 | 0/1 | 0/5 | 0 | 1/6 | 0/0 | 0/0 |
| June 19 | -36 | 0 | 0/1 | 0/1 | 0/0 | 0 | 0/0 | 0/0 | 0/0 |
| July 15-16 | -45 | 4 | 0/1 | 1/1 | 0/0 | 3 | 0/0 | 0/0 | 0/0 |
| Aug. 3-4 | -49 | 5 | 0/0 | 1/1 | 1/1 | 0 | 1/1 | 0/0 | 0/1 |
| Sept. 2-3 | -98 | 3 | 0/1 | 1/1 | 0/0 | 0 | 1/2 | 0/0 | 1/1 |
| Sept. 18-20 | -56 | 5 | 1/1 | 0/1 | 1/3 | 0 | 1/3 | 0/0 | 1/1 |
| Sept. 21 | -24 | 2 | 0/1 | 1/1 | 0/1 | 0 | 0/3 | 0/0 | 1/1 |
| Sept. 21-22 | -30 | 2 | 1/1 | 1/1 | 0/0 | 0 | 0/1 | 0/0 | 0/1 |
| Oct. 1-2 | -98 | 7 | 1/1 | 1/1 | 1/1 | 0 | 1/2 | 0/0 | 0/1 |
| Oct. 10-12 | -130 | 4 | 1/1 | 1/1 | 1/2 | 1 | 0/2 | 0/0 | 0/1 |
| Nov. 7-8 | -110 | 6 | 1/1 | 1/1 | 1/3 | 2 | 0/1 | 0/0 | 0/1 |
| Nov. 22-23 | -108 | 10 | 1/1 | 1/1 | 1/1 | 6 | 0/1 | 0/0 | 1/1 |
| Dec. 10-11 | -60 | 4 | 0/1 | 1/1 | 1/3 | 1 | 0/0 | 0/0 | 1/1 |
| Dec. 30-31 | -77 | 5 | 0/1 | 1/1 | 1/1 | 3 | 0/0 | 0/1 | 0/1 |
| 1998 | | | | | | | | | |
| Jan. 7-8 | -83 | 10 | 0/1 | 1/1 | 4/6 | 0 | 0/1 | 0/0 | 0/1 |
| Feb. 4-5 | -34 | 2 | 1/1 | 0/1 | 0/0 | 0 | 0/6 | 0/0 | 0/1 |
| Feb. 17-18 | -100 | 5 | 0/1 | 1/1 | 1/1 | 1 | 0/0 | 1/1 | 0/1 |
| Mar. 4-5 | -36 | 4 | 1/1 | 0/1 | 2/6 | 1 | 0/0 | 0/0 | 0/1 |
| May 2-3 | -85 | 12 | 1/1 | 1/1 | 0/3 | 8 | 0/1 | 1/1 | 0/0 |
| May 4-5 | -205 | 7 | 1/1 | 0/1 | 1/3 | 1 | 0/0 | 0/0 | 0/1 |
| June 2 | -1 | 1 | 0/0 | 1/1 | 0/1 | 0 | 0/0 | 0/0 | 0/1 |
| June 24-25 | -25 | 7 | 0/1 | 1/1 | 0/4 | 5 | 2/6 | 0/1 | 0/1 |
| Sept. 25-26 | -207 | 14 | 1/1 | 1/1 | 1/2 | 6 | 0/1 | 0/0 | 1/1 |
| Oct. 18-20 | -139 | 9 | 0/1 | 1/1 | 6/10 | 2 | 1/2 | 0/0 | 0/1 |
| Nov. 7-8 | -148 | 11 | 0/1 | 1/1 | 7/17 | 2 | 1/4 | 0/0 | 0/0 |
| Nov. 8-10 | -148 | 23 | 1/1 | 0/0 | 10/12 | 11 | 1/4 | 0/0 | 1/1 |
| Nov. 13-14 | -133 | 20 | 1/1 | 1/1 | ? | ? | ? | 0/0 | 1/1 |
| Total | - | 237 | 17/33 | 25/34 | 51/117 | 64 | 17/65 | 3/6 | 8/28 |
| Average | -86 | 7 | | | | | | | |

R_E on the MS dayside up to 5-7 R_E in the tail. In this case the real MP more often occurs to be closer to the Earth than average location predicted by the model.

SW parameters (the plasma pressure P and the IMF B_Z component) were determined for each MP crossing with taking into account the time delay of plasma propagation between two spacecraft. The range of variation of these parameters for MCs under consideration was found to be sufficiently wide: $0.3 < P < 42$ nPa and $-21 < B_Z < 21$ nT. The ex-

isting MP models (Sibeck *et al.*, 1991; Roelof and Sibeck, 1993; Petrinc and Russell, 1996; Kuznetsov and Suvorova, 1997; Shue *et al.*, 1997) have narrower range of variation. The last version of the MP model (Shue *et al.*, 1998) was obtained with usage of higher SW parameters. By this reason we compared the real MP crossings with two models, and figure 8 presents the distance between the measured location and those predicted in the models by Shue *et al.* (1997) (circles) and Shue *et al.* (1998) (diamonds). In this case, positive




Fig. 6. Observed values of the number of activations NA and minimum of D_{st} for magnetic cloud (black circles) and strong magnetic storms (white circles) periods.




Fig. 7. INTERBALL-1 magnetopause crossings during magnetic cloud passages in the (a) XR_{YZ} and (b) YZ planes.




Fig. 8. Comparison of magnetopause crossings with model (Shue *et al.*, 1997) (circles) and (Shue *et al.*, 1998) (diamonds) predictions.

distances correspond to the event when the measured boundary lies inside model predictions (i.e., closer to the Earth). The distance was measured along the normal to the model boundary.

Figure 8 clearly demonstrates that both the models well predict the MP position in the subsolar region (at $X \geq 0$) and worse in the tail ($X < 0$): on the dayside the MP is located by 1-2 R_E closer to the Earth and in the tail the scatter is from -5 to +2 R_E . Our statistics does not allow us to compare quantitatively both models with sufficient reliability. However, the larger scatter of MP crossing with respect to model predictions testifies that the MP motion during MC passage is more complicated than it is predicted by empirical models which were mainly constructed for the conditions of weakly disturbed SW.

Table 3 presents the results of comparison of the MP location with predictions of the model by Shue *et al.* (1998) as well as the comparison of BS location with its average position. Similar statistical models for BS, which take into account the conditions in the interplanetary space, are absent now; by this reason, the real BS crossing was compared with average BS location. However, since the MP is obstacle for SW in forming BS, we plan to take into account the MP motion depending on conditions in the SW and to investigate the correlation between changes of BS and MP location. Now we can only notice that the deviation of BS from average location is approximately the same as that for MP.

The MP shape and motion for MC of January 10-11, 1997 were studied, in particular, by Nikolaeva *et al.* (1998) and Safrankova *et al.* (1998). The obtained results indicate that the change of MS size was accompanied by more compli-

Table 3. Magnetospheric boundary locations.

| Date | Distance* (R_E) | |
|----------|---------------------|---------------|
| | Bow Shock | Magnetopause |
| 1995 | | |
| Oct. 18 | - | -2.4 ... 1.4 |
| Oct. 19 | - | -4.4 ... 0.6 |
| 1996 | | |
| Dec.25 | ~ 5 | |
| 1997 | | |
| Jan. 10 | - | -0.9 ... 1.4 |
| Jan. 11 | - | 0.0 ... 1.5 |
| Feb. 8 | 3 | - |
| Feb. 9 | 2 ... 3 | - |
| Feb. 10 | 4 | -3.2 |
| Feb. 11 | 3 ... 6 | - |
| June 9 | -5 ... -6 | -0.5 |
| July 3 | -2 ... -4 | - |
| July 4 | 2 | - |
| Sept. 03 | 2 | 1.0 ... 2.8 |
| Sept. 18 | 7 | -3.5 |
| Sept. 20 | 7 | - |
| Sept. 21 | 7 | - |
| 1998 | | |
| Feb. 3 | - | -0.9 |
| Feb. 4 | - | -4.7 ... -2.0 |
| Feb. 18 | - | 1.7 |
| Mar. 4 | -3 ... 3 | 3.5 |
| Mar. 5 | 3 ... 4 | 1.0 |
| May 4 | 0 | 1.7 |

* Distance is positive if the boundary is located closer to the Earth than the model boundary

cated deformations than a simple compression when different parts of the MS simultaneously undergo proportional displacement, by surface waves on the boundaries and by oscillation of the tail (Yermolaev *et al.*, 1997a; Nikolaeva *et al.*, 1998). More complicated character of MS compression follows also from observations on October 18-19, 1995, since these data were interpreted as a result of reconnection of magnetic field not in the subsolar region or near the cusp but rather on the MP in the far tail at distances $|X|$ larger than $20 R_E$ (Savin *et al.*, 1997).

5 Magnetosphere state.

As was shown in previous Section, the MC passage to the Earth is accompanied by the displacement of MS boundaries. This implies partially that the place where one physical region of space is usually observed (which is characterized by typical values of plasma and magnetic field parameters) occurs to be occupied by another region which is observed far from this place under normal conditions. Though small displacement of various regions is a rather frequent phenomenon in such a dynamical system as MS, displacements to distance comparable with the size of regions or even greater are

quite rare phenomena. This fact should be taken into account when comparing the parameters of usual magnetosheath, for instance, with those magnetosheath-like plasma which observed in the region of usual plasma sheet observations. On the other hand, such an analysis is very important since it provides the additional information on the dynamics and mechanisms of different MS region formation. We have considered only several examples from large set of various cases of anomalous location of MS regions, and the results presented below can be considered only as a first step in this direction.

Figure 9 shows the dynamic energy spectrograms of ions (the abscissa is time, the ordinate is energy, the color from blue to red indicates increasing value of ion flux) for three successive orbits of INTERBALL-1 during the period of January 6-16, 1997. In this case the data, placed on the same vertical straight line, were obtained approximately at the same satellite coordinates. (Due to annual satellite orbit evolution with respect to the Sun-Earth axis the planes of successive orbits are displaced relative to each other by an angle of $\sim 4^\circ$.) These data were obtained by the CORALL (Yermolaev *et al.*, 1997b) instrument with the help of a sensor oriented perpendicular to the satellite spin axis, i.e., in the plane normal to the Sun-Earth direction.

The upper panel, whose data were obtained before the MC passage, shows at first a hot and low density plasma of the plasma sheet. During interval from 22 UT on January 6 to 02 UT on January 7, when satellite was close to geomagnetic equator ($X \sim -17$ and $Y_{GSM} \sim 16R_E$) the plasma of a low-latitude boundary layer (LLBL) was observed. After this the satellite began to approach the Earth while crossing several times the plasma mantle and the tail lobes and at ~ 23 UT the satellite reached the radiation belt.

Before the MC passage on January 10 the plasma sheet ions (more precisely PSBL ions) were observed. However, at about 01:20 UT the satellite crossed the MP and entered the magnetosheath. Then, from ~ 06 to ~ 20 UT, the instrument recorded both long (for 1-2 h) intervals and short (a few minutes) bursts of plasma sheet with lower density and higher energy than on the previous orbit. The plasma sheet observations were interrupted by satellite enters the lobes which connected with fast tail motion with respect to a rather slowly moving satellite. After MC trailing edge passage at about 01:20 UT on January 11, the satellite from the plasma sheet quickly entered a very dense and hot magnetosheath, then at ~ 02 UT it was in the LLBL (at a rather large distance from geomagnetic equator with $Z_{GSM} \sim 8 - 9R_E$) and then in the plasma sheet.

The plasma sheet parameters observed on January 13-14 had intermediate values between those observed before and during MC passage. The geomagnetic tail continued to displace relative to the satellite.

The dynamic energy spectrograms of electrons, measured by ION instrument (Sauvaud *et al.*, 1997) on subsequent orbits of the INTERBALL-2 satellite, are presented in Figure 10. Before the MC passage in the polar cap (invariant latitude $\lambda > 65^\circ$) the fluxes of electrons had low energy of several tens of electronvolts and too low intensity to be observed. However after the MC TE passage on January 11, 1997, high fluxes of electron with energy 100-300 eV were detected in the polar cap. This interval coincides with the




Fig. 9. The ion energy spectrograms during 3 successive orbits of INTERBALL-1 on January 6-15, 1997.

INTERBALL-1 exit from the plasma sheet into the magnetosheath and LLBL, i.e., the disturbance of distant tail of MS coincided with electron precipitation in the polar cap.

Thus, several features of the magnetosphere and magnetosheath plasma observed during the MC passage can be summarized as follows.

- The observation of very hot (on January 10-11, 1997 (Yermolaev *et al.*, 1997a, 1998)) as well as very cold (on October 19, 1995 (Savin *et al.*, 1997)) plasma in the magnetosheath.

- A very dense ($N \sim 150 \text{ cm}^{-3}$) plasma in the magnetosheath (Nikolaeva *et al.*, 1998) and no correlation between simultaneous values of density in the magnetosheath and in the plasma sheet (Yermolaev *et al.*, 1997a, 1998).

- The observation of LLBL-like plasma at the large distance ($\sim 8R_E$) from the geomagnetic equator (Yermolaev *et al.*, 1997a, 1998).

- The geomagnetic tail oscillations relative to satellite, so that the displacements of some magnetosphere regions are comparable to characteristic size of the regions (Yermolaev *et al.*, 1997a, 1998, 2000a,b).

- The development of disturbances and acceleration of ions and electrons in the plasma sheet, their subsequent injection and precipitation in polar regions of the magnetosphere (Yermolaev *et al.*, 1997a, 2000a,b).

6 Discussion and conclusions.

Presented results on the analysis of magnetic clouds observed on interplanetary spacecraft and INTERBALL-1,2 satellites during August, 1995 - December, 1998 (near minimum of solar cycle), allow us to make several conclusions about the magnetosphere response to these events.

The geoeffectiveness of magnetic clouds depends on the value of parameter variations in the magnetic cloud. For low, medium, or moderately high variations of plasma and magnetic field in the cloud, the magnetosphere response is the same as for similar variations in the interplanetary space in the absence of magnetic clouds, and strongly depends on the interplanetary magnetic field prehistory:

- after prolonged energy transfer to the magnetosphere (at the southward IMF) practically all changes in the solar wind pressure or in the IMF magnitude and orientation can result in auroral activations, substorms and magnetic storms;

- at prolonged northward IMF all changes in magnetic cloud parameters are not geoeffective and do not have significant influence on the state of the magnetosphere and on the geomagnetic field.

Extremely high jumps of parameters in magnetic clouds (mainly near their boundaries: in shocks, at leading and trail-




Fig. 10. The electron energy spectrograms during 3 orbits of INTERBALL-2 on January 10-12, 1997.

ing edges) can result in the unusual behavior of the magnetosphere:

- strong and rather complicated compression and deformation (with large and disproportional displacement of boundaries) of the magnetosphere relative to its usual position;

- large-scale oscillations of geomagnetic tail structures relative to satellite;

- the development of disturbances in the plasma sheet, which result in acceleration of ions and electrons and their injections in the polar cap.

The magnetic clouds resulting in a greater number of polar disturbances like substorms are accompanied, as a rule, by stronger global disturbance like magnetic storms.

Acknowledgements. The key parameters of plasma and magnetic field in the interplanetary space (PIs of experiments K.W.Ogilvie and R.P.Lepping) were obtained from NASA/GSPC via CDWeb, the data on Auroral Oval Indices from the Cluster/Ground-Based Data Center. We thank S.I.Klimov, S.A.Romanov, M.N.Nozdachev and A.A.Skalsky for results of INTERBALL magnetic field measurements, A.O.Fedorov for his participants in CORALL data processing, R.A.Kovrazhkin and Borodkova for help in ION and ELECTRON data preparation, V.I.Prokhorenko for calculations of INTERBALL orbits, S.P.Savin, L.M.Zelenyi, P.E.Eiges, Z.Nemecek, J.Safrankova, and Yu.I.Galperin for assistance and useful discussions. The work was partially supported by the RFBR, grants 97-02-16489, 98-02-16297 and 98-05-64508, and by the INTAS, grants 97-1612 and 99-0078.

References

- Burlaga, L., R. Fitzenreiter, R. Lepping, K. Ogilvie, A. Szabo, A. Lazarus, J. Steinberg, G. Gloeckler, R. Howard, D. Michels, C. Farrugia, R. P. Lin, and D. E. Larson, A magnetic cloud containing prominence material: January 1997, *J.Geophys.Res.* **103**, 277, 1998.
- Crooker, N. U., Solar and heliospheric geoeffective disturbances, *Journal of Atmospheric and Solar-Terrestrial Physics*, **62**, 1071, 2000.
- Gonzalez, W. D., J. A. Joselyn, Y. Kamide, H. W. Kroehl, G. Rostoker, B. T. Tsurutani, and V. M. Vasyliunas, What is a geomagnetic storm?, *J.Geophys.Res.*, **103**, 277, 1998.
- Gonzalez, W. D., B. T. Tsurutani, and A. L. Clua de Gonzalez, Interplanetary origin of geomagnetic storms, *Space Sci. Rev.*, **88**, 529, 1999.
- Gopalswamy N., A. Lara, R. P. Lepping, M. L. Kaiser, D. Berdichevsky, and O. C. St. Cyr, Interplanetary acceleration of coronal mass ejections, *Geophys.Res. Lett.*, **27**, 145, 2000.
- Gosling, J. T., and V. J. Pizzo, Formation and evolution of corotating interaction regions and their three-dimensional structure, *Space Sci. Rev.*, **89**, 21, 1999.
- Gosling J. T., D. J. McComas, J. L. Phillips, and S. J. Bame, Geomagnetic activity associated with Earth passage of interplanetary shock disturbances and coronal mass ejections, *J.Geophys.Res.*, **96**, 7831, 1991.
- Kamide, Y., W. Baumjohann, I. A. Daglis, W. D. Gonzalez, M. Grade, J. A. Joselyn, R. L. McPherron, J. L. Phillips, E. G. D. Reeves, G. Rostoker, A. S. Sharma, H. J. Singer, B. T. Tsurutani, and V. M. Vasyliunas, Current understanding of magnetic storm: Storm-substorm relationships, *J.Geophys.Res.*, **103**, 17705, 1998.
- Kuznetsov, S. N., and A. N. Suvorova, The magnetopause shape in the region of geosynchronous orbit, *Geomagn. Aeronom.*, **37**, 1, 1997.
- Lepping, R. P., M. H. Acuna, L. F. Burlaga, W. M. Farrel, J. A. Slavin, K. H. Schatten, F. Mariani, N. F. Ness, F. M. Neubauer, Y. C. Whang, J. B. Byrnes, R. S. Kennon, P. V. Panetta, J. Scheifele, and E. M. Worley, The WIND magnetic field investigation, *Space Sci.Rev.*, **71**, 207, 1995.
- Nikolaeva, N. S., G. N. Zastenker, M. N. Nozdachev, A. A. Skalsky, N. A. Eismont, J. Safrankova, Z. Nemecek, O. Santolik, J. Steinberg, A. Lazarus, A. Szabo, R. Lepping, J.-H. Shue, J. Borovsky, M. Tomsen, and L. Frank, Position and motion of the magnetopause during arrival of a magnetic cloud to the Earth on January 10 and 11, 1997, *Kosmich.Issled.*, **36**, 563, 1998 (in Russian).
- Ogilvie, K. W., D. J. Chornay, R. J. Fritzenreiter, F. Hunsaker, J. Keller, J. Lobell, G. Miller, J. D. Scudder, E. C. Sittler Jr., R. B. Torbert, D. Bodet, G. Needell, A. J. Lazarus, J. T. Steinberg, J. H. Tappan, A. Mavretic, and E. Gergin, A comprehensive plasma instrument for the WIND spacecraft, *Space Sci.Rev.*, **71**, 55, 1995.
- Petrinec, S. M., and C. T. Russell, Near-Earth magnetotail shape and size as determined from the magnetopause flaring angle, *J.Geophys.Res.*, **101**, 137, 1996.
- Petrukovich A. A., and S. I. Klimov, The use of solar wind measurements for the analysis and prediction of geomagnetic activity, *Kosmich.Issled.*, **38**, 463, 2000 (in Russian).
- Roelof E. C., and D. G. Sibeck, Magnetopause shape as bivariate function of interplanetary magnetic field Bz and solar wind dynamic pressure, *J.Geophys.Res.*, **98**, 21421, 1993.
- Safrankova, J., Z. Nemecek, L. Prech, G. Zastenker, K. I. Paularena, N. Nikolaeva, M. Nozdachev, A. Skalsky, and T. Mukai, The January 10-11, 1997, magnetic cloud: Multipoint measurements, *Geophys.Res. Lett.*, **25**, 2545, 1998.
- Sauvaud J.-A., H. Barthe, C. Aoustin, J. J. Thocaven, J. Rouzaud, E. Penou, R. A. Kovrazhkin, and K. G. Afanasiev, The ION experiment onboard the INTERBALL-AURORA satellite; initial results on velocity dispersed structures in the cleft and inside the auroral oval, *Ann. Geophys.*, **16**, 1056, 1998.
- Savin S., O. Balan, N. Borodkova, E. Budnik, N. Nikolaeva, V. Prokhorenko, T. Pulkkinen, N. Rybjeva, J. Safrankova, I. Sandahl, E. Amata, U. Auster, G. Bellucci, A. Blagau, J. Blecki, J. Buechner, M. Ciobanu, E. Dubinin, Yu. Yermolaev, M. Echim, A. Fedorov, V. Formisano, R. Grard, V. Ivchenko, F. Jiricek, J. Juchniewicz, S. Klimov, V. Korepanov, H. Koskinen, K. Kudela, R. Lundin, V. Lutsenko, O. Marghitsu, Z. Nemecek, B. Nikutowski, M. Nozdachev, S. Orsini, M. Parrot, A. Petrukovich, N. Pissarenko, S. Romanov, J. Rauch, J. Rustenbach, J.-A. Sauvaud, E.T. Sarris, A. Skalsky, J. Smilauer, P. Triska, J. G. Trotignon, J. Vojta, G. Zastenker, L. Zelenyi, Yu. Agafonov, V. Grushin, V. Khrapchenkov, L. Prech, and O. Santolik, INTERBALL magnetotail boundary case studies, *Adv.Space Res.*, **20**, N 4/5, 999, 1997
- Shue J.-H., J. K. Chao, H.C. Fu, C. T. Russell, P. Song, K. K. Kharana, and H. J. Singer, A new functional form to study the solar wind control of the magnetopause size and shape, *J.Geophys.Res.*, **102**, 9497, 1997.
- Shue J.-H., P. Song, C. T. Russell, J. T. Steinberg, J. K. Chao, G. Zastenker, O. L. Vaisberg, S. Kokubun, H. J. Singer, T. R. Detman, and H. Kawano, Magnetopause location under extreme solar wind conditions, *J. Geophys. Res.*, **103**, 17691, 1998.
- Sibeck, D. G., R. E. Lopez, and E. C. Roelof, Solar wind control of the magnetopause shape, location, and motion, *J. Geophys. Res.*, **96**, 5489, 1991.
- Tsurutani B.T., W.D. Gonzalez, A.L.C.Gonzalez, F. Tang, J.K.Arballo, and M.Okada, Interplanetary origin of geomagnetic activity in the declining phase of the solar cycle, *J. Geophys. Res.*, **100**, 21717, 1995
- Webb D.F., Coronal mass ejections: The key to major interplanetary and geomagnetic disturbances, *Rev.Geophys.*, *Suppl.*, 577, 1995.
- Wilson A. (ed.), *Proceedings of the 5th International Conference*

- on Substorms*, ESA SP-443, 2000.
- Yermolaev, Yu.I., G. N. Zastenker, N. L. Borodkova, R. A. Kovrazhkin, K. Kudela, V. N. Lutsenko, N. S. Nikolaeva, Z. Nemecek, M. N. Nozdrachev, J. Safrankova, J.-A. Sauvaud, A. A. Skalsky, and L. M. Zelenyi**, Magnetic cloud event on 6-11 January, 1997: INTERBALL multi-satellite and multi-instrument observations, *Proc. 31st ESLAB Symp.*, ESA SP-415, 155, 1997a.
- Yermolaev, Yu.I., A. O. Fedorov, O. L. Vaisberg, V. M. Balebanov, Yu. A. Obod, R. Jimenez, J. Fleites, L. Llera, and A. N. Omelchenko**, Ion distribution dynamics near the Earth's bow shock: First measurements with 2-D ion energy spectrometer CORALL on INTERBALL/Tail Probe satellite, *Ann. Geophysicae*, **15**, 533, 1997b
- Yermolaev, Yu.I., G. N. Zastenker, M. N. Nozdrachev, A. A. Skalsky, and L. M. Zelenyi**, Plasma populations in the magnetosphere during the passage of magnetic cloud on 10-11 January, 1997: INTERBALL/Tail Probe observations, *Geophys.Res. Lett.*, **25**, 2565, 1998.
- Yermolaev, Yu. I., G. N. Zastenker, N. L. Borodkova, R. A. Kovrazhkin, N. S. Nikolaeva, M. N. Nozdrachev, S. P. Savin, A. A. Skalsky, L. M. Zelenyi, Z. Nemecek, J. Safrankova, and J.-A.Sauvaud**, Statistic study of magnetosphere response to magnetic clouds: INTERBALL multi-satellite observations, *Phys. Chem.Earth (C)*, **25**, 177, 2000a.
- Yermolaev, Yu. I., G.N.Zastenker, and N.S.Nikolaeva**, The Earth's magnetosphere response to solar wind events according to the INTERBALL project data, *Kosmich.Issled.* **38**, 527, 2000b (in Russian).